

University of Wollongong Research Online

Faculty of Science - Papers (Archive)

Faculty of Science, Medicine and Health

2010

Catalytic solar water splitting inspired by photosynthesis. homogeneous catalysts with a Mechanical ("Machine-like") action

Gerhard F. Swiegers

University of Wollongong, swiegers@uow.edu.au

G Charles Dismukes

Rutgers University

Leone Spiccia

Robin Brimblecombe

Annette Koo

Monash University

See next page for additional authors

Follow this and additional works at: <https://ro.uow.edu.au/scipapers>



Part of the [Life Sciences Commons](#), [Physical Sciences and Mathematics Commons](#), and the [Social and Behavioral Sciences Commons](#)

Recommended Citation

Swiegers, Gerhard F.; Dismukes, G Charles; Spiccia, Leone; Brimblecombe, Robin; Koo, Annette; Chen, Jun; and Wallace, Gordon G.: Catalytic solar water splitting inspired by photosynthesis. homogeneous catalysts with a Mechanical ("Machine-like") action 2010, 1.
<https://ro.uow.edu.au/scipapers/685>

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library: research-pubs@uow.edu.au

Catalytic solar water splitting inspired by photosynthesis. homogeneous catalysts with a Mechanical ("Machine-like") action

Abstract

Chemical reactions may be controlled by either: the minimum threshold energy that must be overcome during collisions between reactant molecules / atoms (the Activation Energy, E_a), or: the rate at which reactant collisions occur (the Collision Frequency, A) (for reactions with low E_a). Reactions of type (2) are governed by the physical, mechanical interaction of the reactants. Such mechanical processes are unusual, but not unknown in molecular catalysts. We examine the catalytic action and macroscopic properties of several abiological mechanical catalysts and show that they display distinct similarities to enzymes in general. An abiological model of the Photosystem II Water Oxidizing Complex that appears to employ a mechanical action has now been found to be a remarkably active and sustained molecular catalyst of water oxidation when illuminated by sunlight. A free-standing Dye-Sensitized Solar Cell that spontaneously splits water into hydrogen and oxygen has been developed using this catalyst.

Keywords

mechanical, machine, like, homogeneous, action, photosynthesis, inspired, splitting, water, solar, catalytic, catalysts

Disciplines

Life Sciences | Physical Sciences and Mathematics | Social and Behavioral Sciences

Publication Details

Swiegers, G. F., Dismukes, G., Spiccia, L., Brimblecombe, R., Koo, A., Chen, J. & Wallace, G. G. (2010). Catalytic solar water splitting inspired by photosynthesis. homogeneous catalysts with a Mechanical ("Machine-like") action. 240th ACS National Meeting and Exposition (p. 1). United States: American Chemical Society.

Authors

Gerhard F. Swiegers, G Charles Dismukes, Leone Spiccia, Robin Brimblecombe, Annette Koo, Jun Chen, and Gordon G. Wallace

Catalytic Solar Water Splitting Inspired by Photosynthesis. Homogeneous Catalysts with a Mechanical (“Machine-Like”) Action

Gerry Swiegers

*Australian Research Council Centre of Excellence for
Electromaterials Science and Intelligent Polymer Research
Institute
University of Wollongong
Wollongong, NSW 2522
Australia*

INTELLIGENT POLYMER
RESEARCH INSTITUTE



Sub-theme: Molecular Machines

- Much interest in developing "*molecular machines*" that drive chemical reactions
- But, what is the "*mechanical action*" that must occur within a molecular catalyst to turn it into a *molecular machine*?

Two General Methods of Inducing Change

- **Energy Gradient (" *Thermodynamics* ")**

Change driven by an overall release of energy


e.g. a ball falling to Earth under gravity

- **Mechanical Interaction (" *Mechanics* ")**

Change driven by a physical collision (action-reaction sequence that plays out over time)

e.g. two billiard balls physically colliding



Chemistry: Collision Theory



What Happens in Catalysis?


STEP (1)
Reactant collision controlled by the
"collision frequency" (A)

STEP (2)
Product formation controlled by the
"activation energy" (E_a)



"Mechanical" Catalysts: H₂ generation






Dynamic proton binding



produces 5 molecules H₂ s⁻¹ catalyst⁻¹ over at least 5 days of continuous operation

-Two dynamic processes (catalyst flexing and proton binding) which are only *synchronized* if the catalyst flexes rapidly about a structure that *complements the transition state*

"Mechanical" Catalysts: O₂ reduction



"Pac-Man" Catalyst

Chem Commun **2007**, 3352

Chem Eur J **2009**, 15, 4746


Some Common Features of “Mechanical” Molecular Catalysts

- (1) Reaction controlled by the *Catalyst-Mediated Collision Frequency* (low *Activation Energy*)
- (2) The maximum catalytic rate depends on the rate of conformational flexing (conformational flexing = the “mechanical impetus”)
- (3) Catalyst typically flexes rapidly about a shape that *complements the transition state*
- (4) Highly efficient and selective form of catalysis (like a machine)
- (5) Michaelis-Menten kinetics

Chem Eur J **2009**, 15, 4746

• These features also found in many enzymes

QUESTION: Are enzymes mechanical catalysts?




Mechanical Catalysis: Methods of Enzymatic, Homogeneous, and Heterogeneous Catalysis


Swiegers, G. F.

John Wiley & Sons, New York, 2008

Can we Mimic an Enzyme?



Water-Oxidizing Complex of
Photosystem II
(PSII-WOC)




Bio-inspired Mn-oxo *Cubane* Model Complex

G. C. Dismukes, Princeton University

Cubane:

- Shape and structure similar to enzyme active site
- Dynamically self-assembles
- Flexible
- Low activation energy for O₂ release (photolytic)

Cubane forms O₂ when illuminated





DeAngelis, Carr

Cubane in Nafion layer on GC electrode

Electrode biased at 1.0 V (vs. Ag/AgCl)


Illuminated at 250-750 nm




Peak turnover frequency:
270 molecules O_2 h^{-1} catalyst $^{-1}$

Total turnovers:
>1000 readily achieved

Cubane⁺ ion-exchanged into Nafion: CV





Proposed Mechanism






Catalyst dis-assembles and re-assembles under turnover conditions

Phys Chem Chem Phys **2009**, 11, 6441
Inorg Chem **2009**, 48, 7269



CV



Water-Splitting “Graetzel” Cell




Water-Splitting “Graetzel” Cell



Uses H₂O as electrolyte – even impure H₂O (seawater).
Eliminates the need for acetonitrile electrolyte which bedevils existing Graetzel cells

J Am Chem Soc **2010**, 132, 2892

- System splits seawater into pure oxygen (no chlorine formation)
- Spin-off company: “*Cube Catalytics LLC*”
(funded by New Energy Ventures, New Jersey)



Acknowledgements

Princeton University / Rutgers University, USA

Prof Chuck Dismukes

Dr Greg Felton

Monash University, Australia

Prof Leone Spiccia

Dr Rob Brimblecombe

Dr Rosalie Hocking

Dr Annette Koo

University of Wollongong, Australia

Prof Gordon Wallace

Dr Jun Chen

Dr Chee Too

CSIRO Molecular and Health Technologies, Australia

Dr Junhua Huang

Funding

Australian Research Council

Australian Academy of Science

University of Wollongong

CSIRO